

BAMF Health Theranostics Center • Molecular Therapy Clinic 109 Michigan Street NW, Suite #200, Grand Rapids, MI 49503 Phone: (888) 870-8998 • Fax: (616) 253-8365 bamfhealth.com

LEQEMBI Order

Revision 01-18-2025

Please Fax completed order to BAMF Health at Fax # 616-253-8365	
PATIENT INFORMATION Referral Sta	tus: 🔄 New Referral 🔄 Updated Order 🗌 Order Renewal
Patient Name:	DOB:
Allergies:	Weight (kg): Height (cm):
ICD-10 Code(s) & Description:	
NOTE: Please send all relevant demographic and insurance ir	
	ase fax copy to BAMF Health) 🔄 NO
REFERRING OFFICE	
	ct Phone #:
	ovider NPI:
	none #: Fax #:
	0.4) Alabaimay's Disease with Late Orest
 Mild cognitive impairment, so stated (ICD-10 code: G31 Alzheimer's Disease with Early Onset (ICD-10 code: G3 	
Other:	Mild dementia due to Alzheimer's Disease
(ICD-10 Code:)	(ICD-10 code: G30.9)
CLINICAL HISTORY – REQUIRED INFORMATION	
Current Medication list	Confirmation of functional impairment w/ validated tool:
Recent provider note(s)	FAQ Score FAST CDR-SB
PET scan or CSF results with amyloid beta confirmation	For Medicare / Medicare Advantage, please include a
Recent MRI of brain (within past year)	copy of the CMS Registry enrollment that includes
Results of cognitive assessment with score:	submission number
(MMSE 22-30,CDR-GS 0.5 or 1)	\Box APoE ε 4 test results (if applicable)
PRE-MEDS FOR 1 st and 2 nd DOSE ONLY	PRE MEDS FOR SUBSEQUENT DOSES
Acetaminophen (Tylenol®) 650 mg PO prior to infusion	Acetaminophen (Tylenol®) 650 mg PO prior to infusion
Decadron 10 mg IVP over 10 minutes prior to infusion	
Please instruct patient to take acetaminophen 650 mg every 6 hours routinely	
for the next 24 hours while at home.	
NURSING ORDERS	
□ Vitals prior to infusion If SBP \ge 150 mmHg, hold treatment and call provider.	
Vitals q15 min x1, end of infusion and prior to discharge	
Patients to be monitored for 1 hr following their first and second infusion	
Call prescriber and/or 911 for any hypersensitivity reactions including but not limited to fever, flu-like symptoms, nausea, vomiting, hypertension, and oxygen desaturation	
THERAPY ADMINISTRATION	
Drug: LEQEMBI (lecanemab-irmb) Dose: 10mg/kg Route: IV over 60 minutes Frequency: Every 2 Weeks Refill: 6 months 1 year Other:	
Date of last infusion if not at BAMF Health: RX Expiration Date:	
FOLLOW UP MR BRAIN w/o contrast (Please use checkboxes below for allowing BAMF Health Nurse navigators to coordinate follow up MRIs at BAMF Health) Rule Out: Amyloid related imaging abnormalities (Brain Edema or Hemorrhage) Other:	
MR Brain w/o contrast (prior to 5 th Lecanemab infusion)	
Order: MR Brain w/o contrast (prior to 5 th Lecanemab infusion)	
MR Brain w/o contrast (prior to 14 th Lecane	
Additional Orders: By signing this order you agree to the following orders, unless otherwise noted.	
 Hold infusion and notify provider if patient reports: Headache, Dizziness, Nausea, Vision Changes, New or Worsening 	
Confusion.	
May initiate IV catheter patency protocol	
 Infusion/allergic reactions may be managed per facility protocol. 	
Provider Signature:	Date: